Cattle Vaccine Comparison Chart

- ●Modified Live Vaccine
- ●Killed Vaccine
- ●Avirulent Live Culture
- ●Approved for Newborn Calves or Pregnant Cows
- ●Safe for newborn calves if dam vaccinated prebreeding according to label
- ●Safe for pregnant cow is vaccinated prebreeding according to label
- ●Safe for pregnant cows or calves nursing pregnant cows if females were vaccinated pre-breeding according to the label

Safe for Newborn Calves	Safe for Pregnant Cows	DO NOT USE in Pregnant Cows/Calves Nursing Pregnant Cows	Respiratory/Respiratory Combinations	*Anthrax*	*Cryptosporidium Parvum*	*IBR*	*BVD 1*	*BVD II*	*PI3*	*BRSV*	*Lepto 5-Way*	*Lepto Hardjo*	*Vibrio (campylobacter)*	*Haemophilus Somnus*	*Pasteurella Multocida*	*Mannheimia Haemolytica*	*E. Coli*	*Coronavirus*	*Bovine Rotavirus*	*Salmonella*	*Clostridia*	*Clostridia B, C & D*	*Clostridium 7-Way*	*Clostridium 8-Way*	*Clostridium 9-Way*	*Cl. Perfringens Type A*	*Tetanus*	*Mycoplasma Bovis*	*Moraxella Bovis (Pinkeye)*	*Moraxella Bovoculi (Pinkeye)*	*Fusobacterium (Foot Rot)*	*Papilomavirus (Warts)*	*Staph Aureus*	*Streptococcus (Mastitis)*	*Klebsiella pneumoniae*	*Rabies*	Dose in ml	Route of Administration	Slaughter Withdrawal
●	●		Bar Somnus-2P											●	●	●																					2 ml	IM	21 days
	●		Bovi-Sera Serum Antibodies												●	●	●			●																	20 ml	IM, SQ	21 days
●	●		Bovi-Shield Gold 5			●	●	●	●	●																										2 ml	IM, SQ	21 days
●	●		Bovi-Shield Gold BVD				●	●																												2 ml	IM	21 days
●	●		Bovi-Shield Gold FP 5			●	●	●	●	●																										2 ml	IM	21 days
●	●		Bovi-Shield Gold FP 5 L5			●	●	●	●	●	●																									2 ml	IM	21 days
●	●		Bovi-Shield Gold FP 5 L5 HB			●	●	●	●	●	●	●																								2 ml	IM	21 days
●	●		Bovi-Shield Gold FP 5 VL5			●	●	●	●	●	●		●																							2 ml	IM	21 days
●	●		Bovi-Shield Gold FP 5 VL5 HB			●	●	●	●	●	●	●	●																							2 ml	IM	21 days
●	●		Bovi-Shield Gold IBR-BVD			●	●	●																												2 ml	IM, SQ	21 days
●	●		Bovi-Shield Gold One Shot			●	●	●	●	●							●																				2 ml	SQ	21 days
	●		CattleMaster 4+VL5			●	●		●	●	●		●																							5 ml	IM	21 days
●	●		CattleMaster Gold FP 5			●	●	●	●	●																										2 ml	SQ	21 days
●	●		CattleMaster Gold FP 5 L5			●	●	●	●	●	●																									5 ml	SQ	21 days
	●		Elite 4-HS			●	●		●	●				●																						5 ml	IM	21 days
	●		Elite 9-HS			●	●		●	●	●			●																						5 ml	IM	21 days
	●		Express 5			●	●	●	●	●																										2 ml	SQ	21 days
	●		Express 5-HS			●	●	●	●	●					●																						2 ml	SQ	21 days
	●		Express FP 3-VL5			●	●	●			●		●																							2 ml	IM, SQ	21 days
	●		Express FP 5-VL5			●	●	●	●	●	●		●																							2 ml	IM, SQ	21 days
	●		Express FP 10			●	●	●	●	●	●	●																								2 ml	IM, SQ	21 days
●	●		Inforce 3			●			●	●																										2 ml	IN	21 days
	●		Mannheimia Haemolytica												●	●																				2 ml	SQ	21 days
	●		Master Guard 10 HB			●	●	●	●	●	●	●																								3 mL	IM, SQ	21 days
●	●		Nasalgen IP			●			●																											2 ml	IN	21 days
		●	Nuplura PH													●																			2 ml	SQ	21 days
●	●		Once PMH IN												●	●																				2 ml	IN	21 days
●	●		Once PMH SQ												●	●																				2 ml	SQ	21 days
●	●		One Shot													●																				2 ml	SQ	21 days
●	●		One Shot BVD				●	●								●																				2 ml	SQ	21 days
		●	Pregguard Gold FP 10			●	●	●	●		●		●																							2 ml	IM	21 days
	●		Presponse HM												●	●																				2 ml	IM	21 days
	●		Presponse SQ													●																				2 ml	SQ	21 days
	●		Pulmo-Guard PH-M												●	●																				2 ml	SQ	21 days
	●		Pulmo-Guard PHM-1												●	●																				2 ml	SQ	60 days
●	●		Pyramid 10			●	●	●	●	●	●																									2 ml	SQ	21 days
	●		Pyramid 5			●	●	●	●	●																										2 ml	SQ	21 days
	●		Pyramid 5 + Presponse SQ			●	●	●	●	●							●																				2 ml	SQ	21 days
●			Somnu Shield												●																						2 ml	IM, SQ	21 days
●	●		Somubac												●																						2 ml	IM, SQ	21 days
	●		Titanium 5			●	●	●	●	●																										2 ml	SQ	21 days
	●		Titanium 5 + PH-M			●	●	●	●	●						●	●																				2 ml	SQ	21 days
	●		Titanium 5 L5 HB			●	●	●	●	●	●	●																									2 ml	IM, SQ	21 days
	●		Triangle 10 HB			●	●	●	●	●	●	●																									5 ml	IM, SQ	21 days
	●		Triangle 4+PH-K			●	●		●	●							●																				5 ml	SQ	21 days
	●		Triangle 5			●	●	●	●	●																											2 ml	IM, SQ	21 days
	●		Triangle 9+PH-K			●	●		●	●	●						●																				5 ml	SQ	21 days
●	●		TSV-2			●			●																											2 ml	IN	21 days
	●		Vira Shield 4+VL5 HB			●	●	●			●	●	●																								5 ml	SQ	60 days

pbsanimalhealth Serving you since 1941
Dairy Health USA. HORSE HEALTH USA

Disclaimer: The information shown here has been gathered from many sources and with great care. Recognizing, however, the possibility of human and typographical errors in transposing such a large amount of technical information, PBS Animal Health hereby disclaims responsibility.

Last Updated 3/15/2019

Cattle Vaccine Comparison Chart

- ●Modified Live Vaccine
- ●Killed Vaccine
- ●Avirulent Live Culture
- ●Approved for Newborn Calves or Pregnant Cows
- ●Safe for newborn calves if dam vaccinated prebreeding according to label
- ●Safe for pregnant cow is vaccinated prebreeding according to label
- ●Safe for pregnant cows or calves nursing pregnant cows if females were vaccinated pre-breeding according to the label

Safe for Newborn Calves	Safe for Pregnant Cows	DO NOT USE in Pregnant Cows/Calves Nursing Pregnant Cows		*Anthrax*	*Cryptosporidium Parvum*	*IBR*	*BVD 1*	*BVD II*	*PI3*	*BRSV*	*Lepto 5-Way*	*Lepto Hardjo*	*Vibrio (campylobacter)*	*Haemophilus Somnus*	*Pasteurella Multocida*	*Mannheimia Haemolytica*	*E. Coli*	*Coronavirus*	*Bovine Rotavirus*	*Salmonella*	*Clostridia*	*Clostridia B, C & D*	*Clostridium 7-Way*	*Clostridium 8-Way*	*Clostridium 9-Way*	*Cl. Perfringens Type A*	*Tetanus*	*Mycoplasma Bovis*	*Moraxella Bovis (Pinkeye)*	*Moraxella Bovoculi (Pinkeye)*	*Fusobacterium (Foot Rot)*	*Papillomavirus (Warts)*	*Staph Aureus*	*Streptococcus (Mastitis)*	*Klebsiella pneumoniae*	*Rabies*	**Dose in ml**	**Route of Administration**	**Slaughter Withdrawal**
	●		Vira Shield 6			●	●	●	●	●																									5 ml	SQ	60 days
	●		Vira Shield 6+L5			●	●	●	●	●	●																									5 ml	SQ	60 days
	●		Vira Shield 6+L5 HB			●	●	●	●	●	●	●																									5 ml	SQ	60 days
	●		Vira Shield 6+L5 HB Somnus			●	●	●	●	●	●	●		●																							5 ml	IM	60 days
	●		Vira Shield 6+L5 Somnus			●	●	●	●	●	●			●																							5 ml	IM	60 days
	●		Vira Shield 6+Somnus			●	●	●	●	●				●																							5 ml	IM	60 days
	●		Vira Shield 6+VL5			●	●	●	●	●	●		●																							5 ml	SQ	60 days
	●		Vira Shield 6+VL5 HB			●	●	●	●	●	●	●	●																									5 ml	SQ	60 days
	●		Vira Shield 6+VL5 HB Somnus			●	●	●	●	●	●	●	●	●																									5 ml	IM	60 days
	●		Vira Shield 6+VL5 Somnus			●	●	●	●	●	●			●																									5 ml	IM	60 days
		●	Vista 5 L5 SQ			●	●	●	●	●	●																											2 ml	SQ	21 days
		●	Vista 5 SQ			●	●	●	●	●																												2 ml	SQ	21 days
		●	Vista 5 VL5 SQ			●	●	●	●	●	●			●																									2 ml	SQ	21 days
		●	Vista Once SQ			●	●	●	●	●							●	●																			2 ml	SQ	21 days
			Mycoplasma																																				
	●		MpB Guard																											●							2 ml	SQ	60 days
	●		Myco-B																											●							2 ml	SQ	60 days
			Salmonella																																				
	●		Endovac - Beef													●	●	●			●																2 ml	IM	21 days
●			EnterVene-d																		●																2 ml	SQ	21 days
●	●		Quatracon-2x													●		●			●																15 ml	SQ	21 days
●	●		Salmonella Dublin-Typhimurium Bacterin																		●																2 ml	SQ	21 days
	●		Salmonella Newport Bacterial Extract																		●																2 ml	SQ	60 days
			Clostridial /Clostridial Combinations																																				
	●		20/20 Vision 7																					●							●						2 ml	SQ	21 days
	●		Alpha-7																					●													2 ml	SQ	21 days
●	●		Alpha-7/MB-1																					●							●						2 ml	SQ	60 days
	●		Bar-Vac 7																					●													5 ml	SQ	21 days
	●		Bar-Vac CD																				●														5 ml	SQ	21 days
	●		Bar-Vac CD/T																				●						●								5 ml	SQ	21 days
	●		Bar-Vac-7/Somnus													●								●													5 ml	SQ	21 days
	●		C & D Antitoxin																				●														10 ml	SQ	21 days
	●		Caliber 7																					●													2 ml	SQ	21 days
	●		Cavalry 9																							●											2 ml	SQ	21 days
	●		Clostratox BCD																				●														15 ml	SQ	21 days
	●		Clostridium Perfringens Type A Toxoid																							●											2 ml	SQ	21 days
●	●		Clostridium Perfringens Types C & D Antitoxin																				●														25 ml	SQ	21 days
	●		Covexin 8																						●												5 ml	IM, SQ	21 days
	●		One Shot Ultra 7														●							●													2 ml	SQ	21 days
	●		One Shot Ultra 8														●								●													2 ml	SQ	21 days
	●		Piliguard Pinkeye+7																					●							●						5 ml	SQ	21 days
	●		Ultrabac 7																					●													5 ml	SQ	21 days
	●		Ultrabac 7/Somubac													●								●													5 ml	SQ	21 days
	●		Ultrabac 8																						●												5 ml	SQ	21 days
	●		Ultrabac CD																				●														2 ml	SQ	21 days
	●		Ultrachoice 7																					●													2 ml	SQ	21 days
	●		Ultrachoice 8																						●												2 ml	SQ	21 days
	●		Vision 7 with SPUR																					●													2 ml	SQ	21 days
	●		Vision 7 Somnus with SPUR													●								●													2 ml	SQ	21 days
	●		Vision 8 with SPUR																						●												2 ml	SQ	21 days
	●		Vision CD-T																				●						●								2 ml	SQ	21 days
			Lepto & Vibrio																																				
●			Citadel L5										●																								2 ml	SQ	21 days
	●		Citadel VL5										●		●																						2 ml	SQ	21 days

pbsanimalhealth Serving you since 1941
Dairy Health USA. HORSE HEALTH USA

Disclaimer: The information shown here has been gathered from many sources and with great care. Recognizing, however, the possibility of human and typographical errors in transposing such a large amount of technical information, PBS Animal Health hereby disclaims responsibility.

Last Updated 3/15/2019

Cattle Vaccine Comparison Chart

- ●Modified Live Vaccine
- ●Killed Vaccine
- ●Avirulent Live Culture
- ●Approved for Newborn Calves or Pregnant Cows
- ●Safe for newborn calves if dam vaccinated prebreeding according to label
- ●Safe for pregnant cow is vaccinated prebreeding according to label
- ●Safe for pregnant cows or calves nursing pregnant cows if females were vaccinated pre-breeding according to the label

Safe for Newborn Calves	Safe for Pregnant Cows	DO NOT USE in Pregnant Cows/Calves Nursing Pregnant Cows		*Anthrax*	*Cryptosporidium Parvum*	*IBR*	*BVD 1*	*BVD II*	*PI3*	*BRSV*	*Lepto 5-Way*	*Lepto Hardjo*	*Vibrio (campylobacter)*	*Haemophilus Somnus*	*Pasteurella Multocida*	*Mannheimia Haemolytica*	*E. Coli*	*Coronavirus*	*Bovine Rotavirus*	*Salmonella*	*Clostridia*	*Clostridia B, C & D*	*Clostridium 7-Way*	*Clostridium 8-Way*	*Clostridium 9-Way*	*Cl. Perfringens Type A*	*Tetanus*	*Mycoplasma Bovis*	*Moraxella Bovis (Pinkeye)*	*Moraxella Bovoculi (Pinkeye)*	*Fusobacterium (Foot Rot)*	*Papilomavirus (Warts)*	*Staph Aureus*	*Streptococcus (Mastitis)*	*Klebsiella pneumoniae*	*Rabies*	**Dose in ml**	**Route of Administration**	**Slaughter Withdrawal**
	●		L5 SQ								●																								2 ml	SQ	21 days
	●		Lepto Shield 5								●																								2 ml	IM	21 days
	●		Leptoferm-5								●																								2 ml	IM	21 days
	●		Reprostar L5 HB								●	●																							2 ml	SQ	60 days
	●		Reprostar VL5 HB								●	●	●																						2 ml	IM	60 days
●	●		Spirovac									●																							2 ml	SQ	21 days
	●		Spirovac L5								●	●																							2 ml	IM, SQ	21 days
●	●		Spirovac VL5								●	●	●																						5 ml	IM, SQ	21 days
	●		Staybred VL5								●		●																						2 ml	IM	21 days
	●		Vib Shield Plus L5								●		●																						2 ml	IM	60 days
	●		VL5 SQ								●	●	●																						2 ml	SQ	21 days
			Pinkeye																																			
●	●		Alpha 7/MB-1																					●					●						2 ml	SQ	60 days
●	●		I-Site XP																										●						2 ml	SQ	21 days
●	●		Moraxella Bovoculi Bacterin																													●			2 ml	SQ	21 days
	●		OCU-Guard MB-1																										●						2 ml	SQ	21 days
●	●		Piliguard Pinkeye -1																										●						2 ml	SQ	60 days
	●		Piliguard Pinkeye +7																					●					●						5 ml	SQ	21 days
	●		Pinkeye Shield XT4																										●						2 ml	IM	60 days
	●		Solidbac Pinkeye IR/PR																										●						--	SQ	21 days
			Scours																																			
●			Bar-Guard-99														●																		10 ml	O	21 days
●			Bo-Bac-2X													●	●			●															15 ml	SQ	21 days
●			Bovilis Coronavirus															●																	2 ml	IN	21 days
●			Bovine Ecolizer														●																		10 ml	O	\
●			Bovine Ecolizer+C20														●				●														20 ml	O	\
	●		Bovine Pili Shield+C														●				●														1 ml	IM	60 days
●	●		Calf-Guard															●	●																3 ml	O	21 days
●			First Defense Boluses														●	●																	1	O	0 days
●			First Defense Technology Concentrate														●	●																	1	O	0 days
●			First Defense Technology Gel														●	●																	30 ml	O	0 days
●			First Defense Tri-Shield														●	●	●																30 ml	O	0 days
	●		Guardian														●	●	●		●														2 ml	SQ	60 days
	●		Scour Bos 4															●	●																2 ml	IM	60 days
	●		Scour Bos 9														●	●	●		●														2 ml	IM	60 days
	●		Scourguard 4K														●	●	●																2 ml	IM	21 days
	●		Scourguard 4KC														●	●	●		●														2 ml	IM	21 days
			Mastitis Prevention																																			
	●		Bovilis J-5														●																		5 ml	SQ	60 days
	●		Endovac Dairy													●	●			●															2 ml	IM	21 days
	●		Enviracor J-5														●																		5 ml	SQ	60 days
	●		J-Vac														●			●															2 ml	IM, SQ	21 days
	●		KLEBVax SRP Klebsiellla Pneumoniae Bacterial Extract																															●	2 ml	SQ	60 days
	●		Lysigin																														●		5 ml	IM	21 days
			Footrot & Warts																																			
	●		Fusoguard																											●					2 ml	SQ	21 days
	●		Serpens Species Bacterin																												●				5 ml	SQ	21 days
	●		Wart Vaccine																												●				10 ml	SQ	21 days

pbsanimalhealth Serving you since 1941
Dairy Health USA. HORSE HEALTH USA

Disclaimer: The information shown here has been gathered from many sources and with great care. Recognizing, however, the possibility of human and typographical errors in transposing such a large amount of technical information, PBS Animal Health hereby disclaims responsibility.

Last Updated 3/15/2019

Cattle Vaccine Comparison Chart

Safe for Newborn Calves	Safe for Pregnant Cows	DO NOT USE in Pregnant Cows/Calves Nursing Pregnant Cows	•Modified Live Vaccine •Killed Vaccine •Avirulent Live Culture •Approved for Newborn Calves or Pregnant Cows •Safe for newborn calves if dam vaccinated prebreeding according to label •Safe for pregnant cow is vaccinated prebreeding according to label •Safe for pregnant cows or calves nursing pregnant cows if females were vaccinated pre-breeding according to the label	*Anthrax*	*Cryptosporidium Parvum*	*IBR*	*BVD 1*	*BVD II*	*PI3*	*BRSV*	*Lepto 5-Way*	*Lepto Hardjo*	*Vibrio (campylobacter)*	*Haemophilus Somnus*	*Pasteurella Multocida*	*Mannheimia Haemolytica*	*E. Coli*	*Coronavirus*	*Bovine Rotavirus*	*Salmonella*	*Clostridia*	*Clostridia B, C & D*	*Clostridium 7-Way*	*Clostridium 8-Way*	*Clostridium 9-Way*	*Cl. Perfringens Type A*	*Tetanus*	*Mycoplasma Bovis*	*Moraxella Bovis (Pinkeye)*	*Moraxella Bovoculi (Pinkeye)*	*Fusobacterium (Foot Rot)*	*Papilomavirus (Warts)*	*Staph Aureus*	*Streptococcus (Mastitis)*	*Klebsiella pneumoniae*	*Rabies*	**Dose in ml**	**Route of Administration**	**Slaughter Withdrawal**
			Tetanus																																				
	●		Tetanus Antitoxin																									●								5 ml	IM, SQ	21 days
	●		Tetanus Toxoid																									●								1 ml	IM	21 days
			Rabies																																			
	●		Imrab 3																																●	2 ml	IM, SQ	21 days
	●		IMRAB Large Animal																																●	2 ml	IM, SQ	21 days
	●		Nobivac 3-Rabies																																●	2 ml	IM	21 days

pbsanimalhealth Serving you since 1941
Dairy Health USA. HORSE HEALTH USA

Disclaimer: The information shown here has been gathered from many sources and with great care. Recognizing, however, the possibility of human and typographical errors in transposing such a large amount of technical information, PBS Animal Health hereby disclaims responsibility.

Last Updated 3/15/2019