

Cattle Vaccine Comparison Chart

Safe for Newborn Calves	Safe for Pregnant Cows	DO NOT USE in Pregnant Cows/Calves Nursing Pregnant Cows																	Dose in ml	Route of Administration	Slaughter Withdrawal															
			IBR	BVD 1	BVD II	PI3	BRSV	Lepto 5-Way	Lepto Hardjo	Vibrio (campylobacter)	Haemophilus Somnus	Pasteurella Multocida	Mannheimia Haemolytica	E. Coli	Coronavirus	Bovine Rotavirus	Salmonella	Clostridia				Clostridia B, C & D	Clostridium 7-Way	Clostridium 8-Way	Clostridium 9-Way	Cl. Perfringens Type A	Tetanus	Mycoplasma Bovis	Moraxella Bovis (Pinkeye)	Moraxella Bovoculi (Pinkeye)	Fusobacterium (Foot Rot)	Papillomavirus (Warts)	Staph. Aureus	Klebsiella Pneumoniae	Rabies	
	♥	Titanium 5 L5 HB	L	L	L	L	L																											2 ml	SQ	21 days
♦		Triangle 4+PH-K	X	X		X	X																											5 ml	SQ	21 days
♦		Triangle 5	X	X	X	X	X																											2 ml	IM, SQ	21 days
♦		Triangle 10 HB	X	X	X	X	X	X	X																									5 ml	IM, SQ	21 days
♦		Vira Shield 4+VL5 HB	X	X	X			X	X	X																								5 ml	SQ	60 days
♦		Vira Shield 6	X	X	X	X	X																											5 ml	SQ	60 days
♦		Vira Shield 6+L5	X	X	X	X	X	X																										5 ml	SQ	60 days
♦		Vira Shield 6+L5 HB	X	X	X	X	X	X	X																									5 ml	SQ	60 days
♦		Vira Shield 6+L5 HB Somnus	X	X	X	X	X	X	X		X																							5 ml	IM	60 days
♦		Vira Shield 6+Somnus	X	X	X	X	X				X																							5 ml	IM	60 days
♦		Vira Shield 6+VL5	X	X	X	X	X	X		X																								5 ml	SQ	60 days
♦		Vira Shield 6+VL5 HB	X	X	X	X	X	X	X	X																								5 ml	SQ	60 days
♦		Vira Shield 6+VL5 HB Somnus	X	X	X	X	X	X	X	X	X																							5 ml	IM	60 days
♦		Vira Shield 6+VL5 Somnus	X	X	X	X	X	X	X	X	X																							5 ml	IM	60 days
	√	Vista 5 L5 SQ	L	L	L	L	L	X	X																								2 ml	SQ	21 days	
	√	Vista 5 SQ	L	L	L	L	L																										2 ml	SQ	21 days	
	√	Vista 5 VL5 SQ	L	L	L	L	L	X	X	X																							2 ml	SQ	21 days	
	√	Vista Once SQ	L	L	L	L	L																										2 ml	SQ	21 days	
Salmonella																																				
♦		EnterVene-d																L															2 ml	SQ	21 days	
♦	♦	Salmonella Dublin-Typhimurium Bacterin																X																2 ml	SQ	21 days
Clostridial /Clostridial Combinations																																				
♦		20/20 Vision 7 with SPUR																		X														2 ml	SQ	21 days
♦		Alpha-7																		X														2 ml	SQ	60 days
♦	♦	Alpha-7/MB-1																		X														2 ml	SQ	60 days
♦		Bar-Vac 7																		X														5 ml	SQ	21 days
♦		Bar-Vac CD																		X														5 ml	SQ	21 days
♦		Bar-Vac CD/T																		X														5 ml	SQ	21 days
♦		Bar-Vac-7/Somnus																		X														5 ml	SQ	21 days
♦		Caliber 7																		X														2 ml	SQ	21 days
♦		Cavalry 9																																2 ml	SQ	21 days
♦		Clostratox BCD																		X														15 ml	SQ	21 days
♦		Clostridium Perfringens Type A Toxoid																																2 ml	SQ	21 days
♦	♦	Clostridium Perfringens Types C & D Antitoxin																		X														25 ml	SQ	21 days
♦		Covexin 8																																5 ml	IM, SQ	21 days
♦		One Shot Ultra 7																		X														2 ml	SQ	21 days
♦		One Shot Ultra 8																		X														2 ml	SQ	21 days
♦		Piliguard Pinkeye+7																		X														5 ml	SQ	21 days
♦		Ultrabac 7																		X														5 ml	SQ	21 days
♦		Ultrabac 7/Somubac																		X														5 ml	SQ	21 days
♦		Ultrabac 8																																5 ml	SQ	21 days
♦		Ultrabac CD																		X														2 ml	SQ	21 days
♦		Ultrachoice 7																		X														2 ml	SQ	21 days
♦		Ultrachoice 8																																2 ml	SQ	21 days

Cattle Vaccine Comparison Chart

Safe for Newborn Calves	Safe for Pregnant Cows	DO NOT USE in Pregnant Cows/Calves Nursing Pregnant Cows	IBR	BVD 1	BVD II	PI3	BRSV	Lepto 5-Way	Lepto Hardjo	Vibrio (campylobacter)	Haemophilus Somnus	Pasteurella Multocida	Mannheimia Haemolytica	E. Coli	Coronavirus	Bovine Rotavirus	Salmonella	Clostridia	Clostridia B, C & D	Clostridium 7-Way	Clostridium 8-Way	Clostridium 9-Way	Cl. Perfringens Type A	Tetanus	Mycoplasma Bovis	Moraxella Bovis (Pinkeye)	Moraxella Bovoculi (Pinkeye)	Fusobacterium (Foot Rot)	Papillomavirus (Warts)	Staph. Aureus	Klebsiella Pneumoniae	Rabies	Dose in ml	Route of Administration	Slaughter Withdrawal		
◆		Endovac Dairy L Modified Live Vaccine X Killed Vaccine C Avirulent Live Culture ◆ Approved for Newborn Calves or Pregnant Cows ◆ Safe for newborn calves if dam vaccinated prebreeding according to label • Safe for pregnant cow if vaccinated prebreeding according to label ◆ Safe for pregnant cows or calves nursing pregnant cows if females were vaccinated prebreeding according to label																																	2 ml	IM	21 days
◆		Enviracor J-5												X																				5 ml	SQ	60 days	
◆		J-Vac												X																				2 ml	IM, SQ	21 days	
◆		KLEBVax SRP Klebsiella Pneumoniae Bacterial Extract																												X			2 ml	SQ	60 days		
◆		Lysigin																											X				5 ml	IM	21 days		
Footrot & Warts																																					
◆		Fusoguard																									X						2 ml	SQ	21 days		
◆		Serpens Species Bacterin																										X					5 ml	SQ	21 days		
◆		Wart Vaccine																										X					10 ml	SQ	21 days		
Tetanus																																					
◆		Tetanus Antitoxin																							X									5 ml	IM, SQ	21 days	
◆		Tetanus Toxoid																						X										1 ml	IM	21 days	
Rabies																																					
◆		Imrab 3																												X				2 ml	IM, SQ	21 days	
◆		IMRAB Large Animal																												X				2 ml	IM, SQ	21 days	
◆		Nobivac 3-Rabies																												X				2 ml	IM	21 days	