

Sheep/Goat Vaccine Comparison Chart

L Modified Live Vaccine
X Killed Vaccine
C Avirulent Live Culture

| | Clostridia | Clostridium 7-way | Clostridium 8-way | Clostridium 9-way | E. Coli | Mannheimia haemolytica | Pasteurella multocida | Salmonella typhimurium | Ovine Ecthyma | Rabies | Tetanus | Campylobacter fetus | Caseous lymphadenitis | Chlamydia psittaci | Dosage | Route of Administration | Slaughter Withdrawal |
|--|------------|-------------------|-------------------|-------------------|---------|------------------------|-----------------------|------------------------|---------------|--------|---------|---------------------|-----------------------|--------------------|--------|-------------------------|----------------------|
| Bar-Vac 7 | | X | | | | | | | | | | | | | 2.5 ml | SQ | 21 days |
| Bar-Vac CD | X | | | | | | | | | | | | | | 2 ml | SQ | 21 days |
| Bar-Vac CD/T | X | | | | | | | | | | X | | | | 2 ml | SQ | 21 days |
| Bovi-Sera Serum Antibodies | | | | | X | X | X | X | | | | | | | 10 ml | IM, SQ | 21 days |
| Caliber 7 | | X | | | | | | | | | | | | | 2 ml | IM | 21 days |
| Campylobacter Fetus Bacterin | | | | | | | | | | | | X | | | 5 ml | SQ | 21 days |
| Case-Bac | | | | | | | | | | | | | X | | 2 ml | SQ | 21 days |
| Chlamydia Psittaci Bacterin | | | | | | | | | | | | | | X | 2 ml | SQ | 60 days |
| Clostratox BCD | X | | | | | | | | | | | | | | 10 ml | SQ | 21 days |
| Clostridium Perfringens Types C & D Antitoxi | X | | | | | | | | | | | | | | 5 ml | SQ | 21 days |
| Covexin 8 | | | X | | | | | | | | | | | | 5 ml | SQ | 21 days |
| Essential 3+T | X | | | | | | | | | | X | | | | 2 ml | IM, SQ | 21 days |
| GoatVac C.D. -T | X | | | | | | | | | | X | | | | 2 ml | IM, SQ | 21 days |
| Imrab 3 | | | | | | | | | | X | | | | | 2 ml | IM, SQ | 21 days |
| IMRAB Large Animal | | | | | | | | | | X | | | | | 2 ml | IM, SQ | 21 days |
| Mannheimia Haemolytica | | | | | | X | | | | | | | | | 2 ml | SQ | 21 days |
| Ovine Ecthyma (Colorado Serum Company) | | | | | | | | | L | | | | | | | S | 21 days |
| Ovine Ecthyma (Texas Vet Lab) | | | | | | | | | L | | | | | | | S | 21 days |
| Tetanus Antitoxin | | | | | | | | | | | X | | | | 5 ml | IM, SQ | 21 days |
| Tetanus Toxoid Concentrate | | | | | | | | | | | X | | | | 1 ml | IM | 21 days |
| Ultrabac 7 | | X | | | | | | | | | | | | | 2.5 ml | SQ | 21 days |
| Ultrabac 8 | | | X | | | | | | | | | | | | 2.5 ml | SQ | 21 days |
| Ultrabac CD | X | | | | | | | | | | | | | | 1 ml | IM, SQ | 21 days |
| UltraChoice 7 | | X | | | | | | | | | | | | | 1 ml | SQ | 21 days |
| UltraChoice 8 | | | X | | | | | | | | | | | | 1 ml | SQ | 21 days |
| Vision 7 with SPUR | | X | | | | | | | | | | | | | 2 ml | SQ | 21 days |
| Vision 8 with SPUR | | | X | | | | | | | | | | | | 2 ml | SQ | 21 days |
| Vision CD-T with SPUR | X | | | | | | | | | | X | | | | 2 ml | SQ | 21 days |