

Cattle Vaccine Comparison Chart

Safe for Newborn Calves	Safe for Pregnant Cows	DO NOT USE in Pregnant Cows/Calves Nursing Pregnant Cows		IBR	BVD 1	BVD II	PI3	BRSV	Lepto 5-Way	Lepto Hardjo	Vibrio (campylobacter)	Haemophilus Somnus	Pasteurella Multocida	Mannheimia Haemolytica	E. Coli	Coronavirus	Bovine Rotavirus	Salmonella	Clostridia	Clostridia B, C & D	Clostridium 7-Way	Clostridium 8-Way	Clostridium 9-Way	Cl. Perfringens Type A	Tetanus	Mycoplasma Bovis	Moraxella Bovis (Pinkeye)	Moraxella Bovoculi (Pinkeye)	Fusobacterium (Foot Rot)	Papillomavirus (Warts)	Staph Aureus	Klebsiella Pneumoniae	Rabies	Dose	Route of Administration	Slaughter Withdrawal				
Respiratory/Respiratory Combinations																																								
◆	◆		Bar Somnus-2P									X	X	X																					2 ml	IM	21 days			
	◆		Bar-Vac-7/Somnus									X									X															5 ml	SQ	21 days		
◆	◆		Bovi-Sera Serum Antibodies									X	X	X				X																	20 ml	IM, SQ	21 days			
♥	●		Bovi-Shield Gold 5	L	L	L	L	L																											2 ml	IM, SQ	21 days			
♥	●		Bovi-Shield Gold BVD		L	L																														2 ml	IM, SQ	21 days		
♥	●		Bovi-Shield Gold FP 5	L	L	L	L	L																												2 ml	IM, SQ	21 days		
♥	●		Bovi-Shield Gold FP 5 L5	L	L	L	L	L	X																											2 ml	IM	21 days		
♥	●		Bovi-Shield Gold FP 5 L5 HB	L	L	L	L	L	X	X																										2 ml	IM, SQ	21 days		
♥	●		Bovi-Shield Gold FP 5 VL5	L	L	L	L	L	X		X																									2 ml	IM	21 days		
♥	●		Bovi-Shield Gold IBR-BVD	L	L	L																														2 ml	IM, SQ	21 days		
♥	●		Bovi-Shield Gold One Shot	L	L	L	L	L						X																						2 ml	SQ	21 days		
◆	◆		Bovilis Nasalgen 3	L			L	L																												2 ml	IN	21 days		
◆	◆		Bovilis Nasalgen 3-PMH	L			L	L						C	C																					2 ml	IN	21 days		
◆			Bovilis Vista BVD CFP		L	L																														2 ml	SQ	21 days		
	◆		CattleMaster 4+VL5	L	X		L	L	X		X																										5 ml	IM	21 days	
◆	◆		CattleMaster Gold FP 5	L	X	X	L	L																													2 ml	SQ	21 days	
◆	◆		CattleMaster Gold FP 5 L5	L	X	X	L	L	X																												5 ml	SQ	21 days	
	♥		Express 5	L	L	L	L	L																													2 ml	SQ	21 days	
	♥		Express 5-HS	L	L	L	L	L				X																									2 ml	SQ	21 days	
	♥		Express FP 5-VL5	L	L	L	L	L	X	X	X																										2 ml	IM, SQ	21 days	
	♥		Express FP 10	L	L	L	L	L	X	X																											2 ml	IM, SQ	21 days	
◆	◆		Inforce 3	L			L	L																													2 ml	IN	21 days	
	◆		Mannheimia Haemolytica										X	X																							2 ml	SQ	21 days	
	◆		Master Guard 10 HB	X	X	X	L	L	X	X																											3 ml	IM, SQ	21 days	
	◆		Myco-B One Dose																								X										2 ml	SQ	60 days	
◆	◆		Nasalgen IP	L			L																														2 ml	IN	21 days	
			Nuplura PH											X																							2 ml	SQ	60 days	
		√	Nuplura PH +3	L	L	L								X																							2 ml	SQ	60 days	
		√	Nuplura PH +BVD		L	L								X																							2 ml	SQ	60 days	
◆	◆		Once PMH IN											C	C																						2 ml	IN	21 days	
◆	◆		Once PMH SQ											C	C																						2 ml	SQ	21 days	
◆	◆		One Shot											X																							2 ml	SQ	21 days	
♥	●		One Shot BVD		L	L								L																							2 ml	SQ	21 days	
		√	PregGuard Gold FP 10	L	L	L	L		X		X																										2 ml	IM	21 days	
	◆		Presponse HM										X	X																							2 ml	IM	21 days	
	◆		Presponse SQ											X																								2 ml	SQ	21 days
	◆		Pulmo-Guard PH-M											X	X																							2 ml	SQ	21 days
	◆		Pulmo-Guard PHM-1											X	X																							2 ml	SQ	60 days
	●		Pyramid 5	L	L	L	L	L																														2 ml	SQ	21 days
	●		Pyramid 5 + Presponse SQ	L	L	L	L	L						X																								2 ml	SQ	21 days
♥	●		Pyramid 10	L	L	L	L	L	X																													2 ml	IM, SQ	21 days
			RESPIVax 3 + PH-M	L	L	L							L	L																								2 ml	SQ	21 days
			RESPIVax 5	L	L	L	L	L																														2 ml	SQ	21 days
			RESPIVax 5 + PH-M	L	L	L	L	L					L	L																								2 ml	SQ	21 days

Disclaimer: The information shown here has been gathered from many sources and with great care. Recognizing, however, the possibility of human and typographical errors in transposing such a large amount of technical information, PBS Animal Health hereby disclaims responsibility. Always follow current product label directions to ensure your vaccination program is effective, to prevent residues in slaughtered animals and milk, and to prevent possible harm to your animals.

Cattle Vaccine Comparison Chart

Safe for Newborn Calves	Safe for Pregnant Cows	DO NOT USE in Pregnant Cows/Calves Nursing Pregnant Cows	IBR	BVD 1	BVD II	PI3	BRSV	Lepto 5-Way	Lepto Hardjo	Vibrio (campylobacter)	Haemophilus Somnus	Pasteurella Multocida	Mannheimia Haemolytica	E. Coli	Coronavirus	Bovine Rotavirus	Salmonella	Clostridia	Clostridia B, C & D	Clostridium 7-Way	Clostridium 8-Way	Clostridium 9-Way	Cl. Perfringens Type A	Tetanus	Mycoplasma Bovis	Moraxella Bovis (Pinkeye)	Moraxella Bovoculi (Pinkeye)	Fusobacterium (Foot Rot)	Papillomavirus (Warts)	Staph Aureus	Klebsiella Pneumoniae	Rabies	Dose	Route of Administration	Slaughter Withdrawal
◆		Ultrabac 8																		X													5 ml	SQ	21 days
◆		Ultrabac CD																X															2 ml	SQ	21 days
◆		UltraChoice 7																	X														2 ml	SQ	21 days
◆		UltraChoice 8																		X													2 ml	SQ	21 days
◆		Vision 7 with SPUR																		X													2 ml	SQ	21 days
◆		Vision 7 Somnus with SPUR									X									X													2 ml	SQ	21 days
◆		Vision 8 with SPUR																		X													2 ml	SQ	21 days
◆		Vision CD-T																X						X									2 ml	SQ	21 days
Lepto & Vibrio																																			
■		Citadel L5						X																									2 ml	IM, SQ	21 days
◆		Citadel VL5						X	X																								2 ml	SQ	21 days
◆		L5 SQ						X																									2 ml	SQ	21 days
◆		Lepto Shield 5						X																									2 ml	IM	21 days
◆		Leptoferm-5						X																									2 ml	IM	21 days
◆		ReproSTAR L5 HB						X	X																								2 ml	SQ	60 days
◆		ReproSTAR VL5 HB						X	X	X																							2 ml	IM	60 days
■	◆	Spirovac							X																								2 ml	SQ	21 days
◆		Spirovac L5						X	X																								2 ml	IM, SQ	21 days
■	◆	Spirovac VL5						X	X	X																							5 ml	IM, SQ	21 days
◆		StayBred VL5						X	X																								2 ml	IM	21 days
◆		VL5 SQ						X	X	X																							2 ml	SQ	21 days
◆		Vib Shield Plus L5						X	X																								2 ml	IM	60 days
Pinkeye																																			
◆		20/20 Vision 7 with SPUR																		X						X							2 ml	SQ	21 days
■	◆	Alpha-7/MB-1																	X							X							2 ml	SQ	60 days
◆		I-Site XP																									X						2 ml	SQ	21 days
■	◆	Moraxella Bovoculi Bacterin - AgriLabs																									X						2 ml	SQ	21 days
■	◆	Moraxella Bovoculi Bacterin - Merck																									X						2 ml	SQ	21 days
◆		Ocu-guard MB-1																								X							2 ml	SQ	21 days
■	◆	Piliguard Pinkeye-1 TRIVALENT																								X							2 ml	IM, SQ	60 days
◆		Piliguard Pinkeye +7																		X						X							5 ml	SQ	21 days
◆		Pinkeye Shield XT4																								X							2 ml	IM	60 days
◆		Solidbac Pinkeye IR/PR																								X							--	SQ	21 days
Scours																																			
■		Bovilis Coronavirus													X																		2 ml	IN	21 days
◆		Bovine Pili Shield+C									X						X																1 ml	IM	60 days
■	◆	Calf-Guard														L	L															3 ml	O	21 days	
■		First Defense Boluses													X	X																	1 bol	O	0 days
■		First Defense Dual-Force													X	X																	1 ds	O	0 days
■		First Defense Technology Concentrate													X	X																	1 scp	O	0 days
■		First Defense Tri-Shield													X	X	X																1 ds	O	0 days
◆		Guardian													X	X	X	X															2 ml	SQ	60 days
◆		Scour Bos 4													X	X																	2 ml	IM	60 days
◆		Scour Bos 9													X	X	X	X															2 ml	IM	60 days
◆		ScourGuard 4K													X	X	X																2 ml	IM	21 days
◆		ScourGuard 4KC													X	X	X	X															2 ml	IM	21 days

Disclaimer: The information shown here has been gathered from many sources and with great care. Recognizing, however, the possibility of human and typographical errors in transposing such a large amount of technical information, PBS Animal Health hereby disclaims responsibility. Always follow current product label directions to ensure your vaccination program is effective, to prevent residues in slaughtered animals and milk, and to prevent possible harm to your animals.

